

## HYPERGROUPS ASSOCIATED TO HARMONIC $NA$ GROUPS

BIANCA DI BLASIO

(Received 31 May 2000; revised 13 January 2001)

Communicated by A. H. Dooley

### Abstract

A harmonic  $NA$  group is a suitable solvable extension of a two-step nilpotent Lie group  $N$  of Heisenberg type by  $\mathbb{R}^+$ , which acts on  $N$  by anisotropic dilations. A hypergroup is a locally compact space for which the space of Borel measures has a convolution structure preserving the probability measures and satisfying suitable conditions. We describe a class of hypergroups associated to  $NA$  groups.

2000 *Mathematics subject classification*: primary 43A62, 43A80.

*Keywords and phrases*: hypergroups, harmonic spaces.

### 0. Introduction

A hypergroup is a locally compact space  $X$  for which the space  $M(X)$  of Borel measures on  $X$  is a  $*$ -algebra satisfying suitable conditions (see [11]). Since any symmetric space  $S$  has a hypergroup  $X(S)$  naturally associated to it, Wildberger [12] pointed out the following problem: for what more general Riemannian manifold  $S$  does the hypergroup  $X(S)$  make sense? In this paper we partially answer this question: we introduce new examples of hypergroups associated to a class of harmonic spaces. We recall that a Riemannian manifold is said to be *harmonic* if the volume density depends only on the geodesic distance from a fixed point.

The  $NA$  groups considered in the paper are one-dimensional solvable extensions of two-step nilpotent Lie groups  $N$  of Heisenberg type equipped with a suitable Riemannian metric [7]. The class of these  $NA$  groups includes all noncompact symmetric spaces of rank one (the real hyperbolic spaces fit into this framework as degenerate cases); however most of them are nonsymmetric harmonic spaces [7]. So far this class of manifolds provides the only known examples of nonsymmetric harmonic spaces.

In the paper we briefly describe hypergroups associated to  $NA$  groups and their main properties. We give the explicit expression for the convolution structure, which can be deduced from a complicated formula proved by Flensted-Jensen and Koornwinder [9]. Finally we notice that hypergroups associated to  $NA$  groups of the same dimension and such that the center of the Lie algebra  $\mathfrak{n}$  of  $N$  have the same dimension are all isomorphic.

### 1. Notation

Denote by  $M(X)$  the space of Radon measures on a locally compact space  $X$  and by  $M_1(X)$  the subset of  $M(X)$  of probability measures. The space  $M(X)$  is the dual of  $C_0(X)$ , the space of continuous functions on  $X$  which vanish at infinity. We shall write

$$(f, \nu) = \int_X f(r) d\nu(r) \quad f \in C_0(X), \nu \in M(X).$$

### Hypergroups

**DEFINITION ([11]).** A locally compact space  $X$  is a *hypergroup* if  $M(X)$  is an associative algebra with respect to a convolution  $\star$  satisfying the following conditions:

- (1)  $M_1(X) \star M_1(X) \subset M_1(X)$ ;
- (2) the convolution is separately continuous from  $M_1(X) \times M_1(X)$  in  $M_1(X)$  with respect to the weak\* topology;
- (3) the map  $(r, r') \mapsto \delta_r \star \delta_{r'}$  is continuous from  $X \times X$  in  $M(X)$  with the weak\* topology;
- (4) there exists an element  $e$  in  $X$ , necessarily unique, such that  $\delta_e \star \nu = \nu \star \delta_e = \nu$  for every  $\nu$  in  $M(X)$ ;
- (5) there exists an involutive homomorphism from  $X$  onto  $X$ ,  $r \mapsto \check{r}$ , which extends naturally to  $M(X)$  and satisfies  $(\mu \star \nu)^\check{ } = \check{\nu} \star \check{\mu}$ , for every  $\mu$  and  $\nu$  in  $M(X)$ ;
- (6)  $r = r'$  if and only if  $e \in \text{supp}(\delta_{r'} \star \delta_{\check{r}})$ ;
- (7) for every compact set  $K \subset X$  and for every neighborhood  $V$  of  $K$ , there exists a neighborhood  $U$  of  $e$  such that
  - (a) if  $\text{supp}(\mu) \subset K$  and  $\text{supp}(\nu) \subset U$ , then  $\text{supp}(\mu \star \nu), \text{supp}(\nu \star \mu) \subset V$ ;
  - (b) if  $\text{supp}(\mu) \subset K$  and  $\text{supp}(\nu) \subset V^c$ , then the supports of  $\mu \star \check{\nu}$  and  $\check{\mu} \star \nu$  are disjoint from  $U$ .

A *character* on a commutative hypergroup  $X$  is a nontrivial, continuous, bounded function  $\underline{\chi} : X \mapsto \mathbb{C}$  such that  $(\chi, \delta_r \star \delta_{r'}) = \chi(r)\chi(r')$ , for every  $r, r'$  in  $X$ . If  $\chi(\check{r}) = \overline{\chi(r)}$ , for every  $r$  in  $X$ , then  $\chi$  is an hermitian character.

**Harmonic NA groups** Let  $\mathfrak{n}$  be a two-step real nilpotent Lie algebra, with an inner product  $\langle \cdot, \cdot \rangle_{\mathfrak{n}}$ . Write  $\mathfrak{n}$  as an orthogonal sum  $\mathfrak{n} = \mathfrak{v} \oplus \mathfrak{z}$ , where  $\mathfrak{z}$  is the center of  $\mathfrak{n}$ . For each  $Z$  in  $\mathfrak{z}$ , define the map  $J_Z : \mathfrak{v} \rightarrow \mathfrak{v}$  by the formula

$$\langle J_Z X, X' \rangle_{\mathfrak{n}} = \langle [X, X'], Z \rangle_{\mathfrak{n}} \quad \forall X, X' \in \mathfrak{v}.$$

We say that the Lie algebra  $\mathfrak{n}$  is *H-type* if

$$J_Z^2 = -|Z|^2 I_{\mathfrak{v}} \quad \forall Z \in \mathfrak{z},$$

where  $I_{\mathfrak{v}}$  is the identity on  $\mathfrak{v}$ . The dimension of  $\mathfrak{v}$  is even,  $\dim \mathfrak{v} = 2m$ , say; we denote the dimension of the center  $\mathfrak{z}$  by  $k$ . A connected and simply connected Lie group  $N$  whose Lie algebra is an *H-type algebra* is said to be an *H-type group*.

Let  $NA$  be the semidirect product of the Lie groups  $N$  and  $A = \mathbb{R}^+$  with respect to the action of  $A$  on  $N$  given by the dilations  $(X, Z) \mapsto (a^{1/2}X, aZ)$ . As customary we write  $(X, Z, a)$  for the element  $na = \exp(X + Z)a$ , where  $X$  is in  $\mathfrak{v}$ ,  $Z$  in  $\mathfrak{z}$  and  $a$  in  $A$ . It can easily be checked that the product law in  $NA$  is given by

$$(X, Z, a)(X', Z', a') = \left( X + a^{1/2}X', Z + aZ' + \frac{1}{2}a^{1/2}[X, X'], aa' \right).$$

The inverse of  $(X, Z, a)$  is  $(-a^{-1/2}X, -a^{-1}Z, a^{-1})$ . We denote by  $Q = m + k$  the homogeneous dimension of  $N$ . A left Haar measure on  $NA$  is given by

$$dx = a^{-Q-1} dX dZ da,$$

where  $dX$ ,  $dZ$  and  $da$  are Lebesgue measures on  $\mathfrak{v}$ ,  $\mathfrak{z}$  and  $\mathbb{R}^+$  respectively. We endow  $NA$  with the left-invariant Riemannian structure induced by the following inner product on the Lie algebra  $\mathfrak{n} \oplus \mathbb{R}$  of  $NA$

$$\langle (X, Z, \alpha), (X', Z', \alpha') \rangle = \langle (X, Z), (X', Z') \rangle + \alpha\alpha' \quad X \in \mathfrak{v}, Z \in \mathfrak{z}, \alpha \in \mathbb{R}.$$

The expression of the volume density in geodesic polar coordinates is [1, 7]

$$\Lambda(\rho) = 2^{2Q} \left( \sinh \frac{\rho}{2} \right)^{2m+k} \left( \cosh \frac{\rho}{2} \right)^k d\rho.$$

The fact that  $\Lambda$  depends only on the geodesic distance  $\rho$  from the identity implies that, as a Riemannian manifold,  $NA$  is a harmonic space [7]. Notice that once we have an *H-type algebra*  $\mathfrak{n}$ , the construction of the associated  $NA$  group is straightforward.

We say that a function  $F : NA \rightarrow \mathbb{C}$  is *radial* if  $F(x)$  depends only on the geodesic distance  $|x|$  of the point  $x$  from the identity  $o = (0, 0, 1)$  in  $NA$ .

Given a locally integrable function  $F$  on  $NA$  we denote by  $\mathcal{R}F$  the radial function on  $NA$  given by

$$(1) \quad \mathcal{R}F(x) = \int_{S_r} F(y) d\sigma_r(y) \quad x \in NA, \quad r = |x|,$$

where  $S_r = \{x \in NA : |x| = r\}$  is the geodesic sphere of radius  $r$  centered at the identity and  $\sigma_r$  is the surface measure supported on  $S_r$  normalized so that  $\int_{S_r} d\sigma_r = 1$ . Given a measure  $\mu$  in  $M(NA)$  we denote by  $\mathcal{R}\mu$  the measure defined by the equality  $(\mathcal{R}F, \mu) = (F, \mathcal{R}\mu)$ , for every  $F$  in  $C_0(NA)$ . We say that a measure  $\mu$  in  $M(NA)$  is radial if  $\mathcal{R}\mu = \mu$ .

## 2. Hypergroups associated to $NA$ groups

Let  $X$  be the locally compact space given by the half line  $[0, +\infty)$  endowed with the Euclidean topology. If  $f$  is in  $C_0(X)$  denote by  $F$  the radial function on  $NA$  defined by  $F(x) = f(|x|)$  for every  $x$  in  $NA$ . For  $r$  and  $r'$  in  $X$  let  $\delta_r \star \delta_{r'}$  be the measure on  $X$  defined by the equality

$$(2) \quad (f, \delta_r \star \delta_{r'}) = (F, \sigma_r \star \sigma_r) = \int_{S_{r'}} \int_{S_r} F(xy) d\sigma_r(x) d\sigma_{r'}(y) \quad \forall f \in C_0(X).$$

The convolution  $\star$  naturally extends to  $M(X)$ .

**THEOREM 2.1.** *The locally compact space  $X$  is a hypergroup with respect to the convolution  $\star$  defined by (2) and the involution  $r \mapsto \check{r} = r$  on  $X$ .*

**PROOF.** The proof is a verification of conditions (1)–(7) defining a hypergroup. A detailed proof is given in [3]. □

Therefore we can associate a hypergroup to every  $NA$  group. Our next aim is to write the explicit expression for the convolution  $\star$  defined by (2).

It is easy to see that  $\Lambda(u) du$  is a translation-invariant positive Radon measure on the hypergroup  $X$ .

For any complex number  $\lambda$  denote by  $\Phi_\lambda : NA \rightarrow \mathbb{C}$  the spherical function on  $NA$  associated to the eigenvalue  $-(\lambda^2 + Q^2/4)$  of the Laplace-Beltrami operator on  $NA$ . The spherical function  $\Phi_\lambda$  is bounded if and only if  $|\text{Im } \lambda| \leq Q/2$  [7]. Denote

$$\varphi_\lambda(r) = \Phi_\lambda(x) \quad \text{if } |x| = r.$$

It is easy to see that the set of the hermitian characters of  $X$  is  $\{\varphi_\lambda : \lambda \in \mathbb{R} \cup [-iQ/2, iQ/2]\}$ .

The spherical functions  $\{\Phi_\lambda\}_{\lambda \in \mathbb{C}}$  on NA can be expressed in terms of hypergeometric functions. More precisely we have ([1, 2])

$$\varphi_\lambda(r) = {}_2F_1(Q/2 - i\lambda, Q/2 + i\lambda; (2m + k + 1)/2; -\sinh^2(r/2)),$$

where  ${}_2F_1$  denotes the hypergeometric function according to the notation in [8].

Flensted-Jensen and Koornwinder proved the following equality [9, formulae (4.2) and (4.17)]

$$(3) \quad \varphi_\lambda(r)\varphi_\lambda(t) = \int_0^{+\infty} \varphi_\lambda(u)T(r, t, u)\Lambda(u) du \quad \forall r, t \in [0, +\infty),$$

where for  $u \notin (|r - t|, |r + t|)$

$$T(r, t, u) = 0$$

and for  $|r - t| < u < |r + t|$

$$T(r, t, u) = \frac{\Gamma((2m + k + 1)/2)}{2^{2Q} \sqrt{\pi} \Gamma(m) \Gamma(k/2)} (\sinh(r/2) \sinh(t/2) \sinh(u/2))^{-2m-k+1} \\ \times \int_0^\pi (g_{(r,t,u)}(\theta))_+^{m-1} (\sin \theta)^{k-1} d\theta,$$

with  $g_{(r,t,u)}(\theta) = 1 - (\cosh \frac{r}{2})^2 - (\cosh \frac{t}{2})^2 - (\cosh \frac{u}{2})^2 + 2 \cosh \frac{r}{2} \cosh \frac{t}{2} \cosh \frac{u}{2} \cos \theta$ . Here  $(z)_+ = z$  for  $z > 0$  and  $(z)_+ = 0$  for  $z \leq 0$ .

**THEOREM 2.2.** For every  $r$  and  $t$  in  $X = [0, +\infty)$ , we have

$$d(\delta_r \star \delta_{r'}) (u) = T(r, t, u)d\eta(u) \quad \forall u \in X.$$

**PROOF.** Let  $r$  and  $t$  be in  $X$  and let  $x$  and  $y$  be in NA such that  $|x| = r$  and  $|y| = t$ . In [7] the following relation is proved

$$\mathcal{R}({}_x\Phi_\lambda)(y) = \Phi_\lambda(x)\Phi_\lambda(y) = \varphi_\lambda(r)\varphi_\lambda(t) \quad \forall \lambda \in \mathbb{C},$$

where  ${}_x\Phi_\lambda(y) := \Phi_\lambda(xy)$ . For  $|\text{Im } \lambda| \leq Q/2$ , denote by  $\widehat{F}(\lambda) = \int_{NA} F(x)\Phi_\lambda(x)dx$  the spherical transform of an integrable radial function  $F$  on NA.

Let  $f$  be a  $C^\infty$  function with compact support on  $X$  and define  $F : NA \rightarrow \mathbb{C}$  by  $F(x) = f(|x|)$ , for every  $x$  in NA. By the inversion formula for the spherical

transform on  $NA$  [10] and by the Fubini Theorem we have

$$\begin{aligned}
 (f, \delta_r \star \delta_{r'}) &= \int_{S_r} \int_{S_{r'}} F(xy) d\sigma_r(x) d\sigma_{r'}(y) \\
 &= \int_{S_r} \int_{S_{r'}} c \int_{-\infty}^{+\infty} \widehat{F}(\lambda) \Phi_\lambda(xy) |\mathbf{c}(\lambda)|^{-2} d\lambda d\sigma_r(x) d\sigma_{r'}(y) \\
 &= c \int_{-\infty}^{+\infty} \int_{S_r} \mathcal{R}_x(\Phi_\lambda)(y) d\sigma_r(x) \widehat{F}(\lambda) |\mathbf{c}(\lambda)|^{-2} d\lambda \\
 &= c \int_{-\infty}^{+\infty} \varphi_\lambda(r) \varphi_\lambda(t) \widehat{F}(\lambda) |\mathbf{c}(\lambda)|^{-2} d\lambda \\
 &= c \int_{-\infty}^{+\infty} \int_0^{+\infty} \varphi_\lambda(u) T(r, t, u) \Lambda(u) du \widehat{F}(\lambda) |\mathbf{c}(\lambda)|^{-2} d\lambda \\
 &= \int_0^{+\infty} f(u) T(r, t, u) \Lambda(u) du,
 \end{aligned}$$

where  $c$  is a constant depending only on  $m$  and  $k$  and

$$\mathbf{c}(\lambda) = \frac{2^{Q-2i\lambda} \Gamma(2i\lambda) \Gamma((2m+k+1)/2)}{\Gamma(Q/2+i\lambda) \Gamma((m+1)/2+i\lambda)}. \quad \square$$

It is known that if  $\mathfrak{n} = \mathfrak{v} \oplus \mathfrak{z}$  and  $\mathfrak{n}' = \mathfrak{v}' \oplus \mathfrak{z}'$  are  $H$ -type algebra such that  $\dim \mathfrak{n} = \dim \mathfrak{n}'$  and  $\dim \mathfrak{z} = \dim \mathfrak{z}'$  then  $\mathfrak{n}$  and  $\mathfrak{n}'$  are not necessarily isomorphic (see [5]); however, the associated hypergroups are isomorphic.

**THEOREM 2.3.** *If  $\mathfrak{n} = \mathfrak{v} \oplus \mathfrak{z}$  and  $\mathfrak{n}' = \mathfrak{v}' \oplus \mathfrak{z}'$  are two  $H$ -type algebras such that  $\dim \mathfrak{n} = \dim \mathfrak{n}'$  and  $\dim \mathfrak{z} = \dim \mathfrak{z}'$ , then the hypergroups associated to  $\exp(\mathfrak{n} + \mathbb{R})$  and  $\exp(\mathfrak{n}' + \mathbb{R})$  are isomorphic.*

**PROOF.** Although the result follows by Theorem 2.2, we give here a different proof which does not involve formula (3). From (2) and from the expression of the product law in  $NA$  it is clear that the result follows if we prove that the convolution of radial functions on  $NA$  does not depend on the bracket of the Lie algebra  $\mathfrak{n}$ . Fix a point  $(X'', Z'', a'')$  in  $NA$  and let  $(0, Z', a')$  be in  $NA$  such that  $|(X'', Z'', a'')| = |(0, Z', a')|$ . Let  $F$  and  $\Psi$  be radial functions on  $NA$ . Since  $F * \Psi$  is a radial function on  $NA$ , we have

$$\begin{aligned}
 (F * \Psi)(X'', Z'', a'') &= (F * \Psi)(0, Z', a') \\
 &= \int_{NA} F(X, Z, a) \Psi((X, Z, a)^{-1}(0, Z', a')) a^{-Q-1} dX dZ da. \\
 &= \int_{NA} F(X, Z, a) \Psi(-a^{-1/2}X, a^{-1}(Z' - Z), a'a^{-1}) a^{-Q-1} dX dZ da.
 \end{aligned}$$

We are done since the last expression does not depend on the Lie bracket of  $n$ .  $\square$

**REMARK.** Wildberger [12] noted that there is a hypergroup associated to any Gelfand pair  $(G, K)$ . When  $NA$  is a noncompact symmetric space  $G/K$ , the compact Lie group  $K$  acts transitively on the geodesic spheres of  $G/K$ ; moreover  $(G, K)$  is a Gelfand pair. For a generic  $NA$  harmonic space does not exist a group  $K$  with these properties (see [6]); however it is possible to generalize the theory of Gelfand pairs to this situation [4]. Let us give more information on this topic.

Damek and Ricci [7] introduced the notion of an averaging projection  $\mathcal{R}$  on a Lie group  $S$ , which is an idempotent operator on the space of  $C^\infty$  functions with compact support on  $S$  satisfying suitable conditions. The functions in the image of  $\mathcal{R}$  are called radial. The convolution of radial functions is a radial function and the space  $L^1_{\mathcal{R}}(S)$  of radial integrable functions is a Banach algebra. If the convolution algebra  $L^1_{\mathcal{R}}(S)$  is commutative it plays the same role played by the Banach algebra  $L^1(K \setminus G/K)$  in the theory of Gelfand pairs [4]. Damek and Ricci [7] proved that the operator  $\mathcal{R}$  defined by (1) is an averaging projection on  $NA$  and that the Banach algebra  $L^1_{\mathcal{R}}(NA)$  is commutative.

This point of view is not emphasized in this paper, however we think that it could be interesting to study connections between hypergroups and Lie groups admitting an averaging projection.

## Acknowledgements

Theorem 2.1 and Theorem 2.3 are contained in the Ph.D. Thesis [3] of the author, who is grateful to her advisor Fulvio Ricci for his precious help and encouragement. The author also wishes to thank Jean-Philippe Anker and Norman Wildberger for many useful conversations.

## References

- [1] J.-Ph. Anker, E. Damek and C. Yacoub, ‘Spherical analysis on harmonic  $AN$  groups’, *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* **23** (1996), 643–679.
- [2] F. Astengo, ‘A class of  $L^p$  convolutors on harmonic extensions of  $H$ -type groups’, *J. Lie Theory* **5** (1995), 147–164.
- [3] B. Di Blasio, *Analisi di Fourier sferica su estensioni risolubili di gruppi di Heisenberg generalizzati* (Ph.D. Thesis, Politecnico di Torino, 1997).
- [4] ———, ‘An extension of the theory of Gelfand pairs to radial functions on Lie groups’, *Boll. Un. Mat. Ital. B (7)* **11** (1997), 623–642.
- [5] M. Cowling, A. H. Dooley, A. Korányi and F. Ricci, ‘ $H$ -type groups and Iwasawa decompositions’, *Adv. Math.* **87** (1991), 1–41.

- [6] E. Damek, 'The geometry of a semidirect extension of a Heisenberg type nilpotent group', *Colloq. Math.* **53** (1987), 255–268.
- [7] E. Damek and F. Ricci, 'Harmonic analysis on solvable extensions of  $H$ -type groups', *J. Geom. Anal.* **2** (1992), 213–248.
- [8] A. Erdelyi, W. Magnus, F. Oberhettinger and G. Tricomi, *Higher transcendental functions*, Vol. I (McGraw-Hill, New York, 1953).
- [9] M. Flensted-Jensen and T. Koornwinder, 'The convolution structure for Jacobi function expansions', *Ark. Mat.* **11** (1973), 245–262.
- [10] F. Ricci, 'The spherical transform on harmonic extensions of  $H$ -type groups', *Rend. Sem. Mat. Univ. Politec. Torino* **50** (1992), 381–392.
- [11] R. Spector, 'Aperçu de la théorie des hypergroupes', in: *Analyse harmonique sur les groupes de Lie (Sém. Nancy-Strasbourg, 1973–75)* Lecture Notes in Math. 497 (Springer, Berlin, 1975) pp. 643–673.
- [12] N. J. Wildberger, 'Hypergroups, symmetric spaces, and wrapping maps', in: *Probability measures on groups and related structures, XI, (Oberwolfach, 1994)* (World Sci. Publishing, River Edge, NJ, 1995) pp. 406–425.

Dipartimento di Matematica  
Università degli studi di Roma 'Tor Vergata'  
via della Ricerca scientifica 1  
00133 Roma  
Italy  
e-mail: [diblasio@axp.mat.uniroma2.it](mailto:diblasio@axp.mat.uniroma2.it)