

# ABELIAN GROUPS WHOSE SUBGROUP LATTICE IS THE UNION OF TWO INTERVALS

SIMION BREAZ and GRIGORE CĂLUGĂREANU

(Received 15 February 2003; revised 21 July 2003)

Communicated by E. A. O'Brien

## Abstract

In this note we characterize the abelian groups  $G$  which have two different proper subgroups  $N$  and  $M$  such that the subgroup lattice  $L(G) = [0, M] \cup [N, G]$  is the union of these intervals.

2000 *Mathematics subject classification*: primary 06C99, 20K10, 20K27.

*Keywords and phrases*: subgroup lattice, interval, torsion abelian group, cocyclic groups.

For every subgroup  $H$  of an arbitrary group  $G$ , the interval  $[H, G]$  is a compactly generated (algebraic) sublattice in the subgroup lattice  $L(G)$ .

After 1989, when Tuma [4] showed that *every algebraic lattice is isomorphic to an interval in the subgroup lattice of some group* (improving Whitman's theorem [5]—*every lattice is isomorphic to a sublattice of the subgroup lattice of a group*—as far as possible), an increasing role of intervals, in subgroup lattices of groups, was noticed.

In [1], an arbitrary group  $G$  was called a *BP-group* if it has a proper subgroup  $H$  such that the subgroup lattice  $L(G)$  is the union of the intervals  $[1, H]$  and  $[H, G]$  (that is, any subgroup of  $G$  is either contained in  $H$  or contains  $H$ ). The subgroup  $H$  was called a *breaking point* for the lattice  $L(G)$ . It was pointed out that the abelian BP-groups are the nonsimple cocyclic groups (that is, up to isomorphism,  $\mathbb{Z}(p^k)$  with  $k > 1$  or  $\infty$ ).

Roland Schmidt suggested the study of finite groups which satisfy a weaker condition: groups  $G$  having two proper subgroups  $N$  and  $M$  such that every subgroup  $H$  of  $G$  either contains  $N$  or is contained in  $M$ . In this situation the subgroup lattice  $L(G)$  is again union of two intervals, namely  $[1, M]$  and  $[N, G]$  (such groups appeared in the study of affinities of groups—see for example [3, 9.4.14]—but there are much more examples of this kind).

In this paper, instead of finite groups, we characterize the abelian groups which share this property. Our result is the following:

**THEOREM 1.** *An abelian group  $G$  has two proper subgroups  $N \neq M$  such that the subgroup lattice  $L(G) = [0, M] \cup [N, G]$  if and only if  $G$  is a torsion group with a primary component  $G_p \cong \mathbb{Z}(p^n) \oplus B$ ,  $n \in \mathbb{N}^* \cup \{\infty\}$  such that  $p^l B = 0$  holds for a nonnegative integer  $l < n$ .*

Additive notation is used and from now on, ‘group’ means ‘abelian group’,  $\mathbb{N}$  denotes the set of all nonnegative integers,  $\mathbb{P}$  denotes the set of all prime numbers and standard interval notation is used. We denote by  $h_p(b)$  the  $p$ -height of  $b$ .

We first mention the following simple necessary condition:  *$N$  must be cyclic.* Indeed, take  $x \in G \setminus M$ . Then  $\langle x \rangle \in [0, M]$  being not possible,  $\langle x \rangle \in [N, G]$  or  $N \leq \langle x \rangle$ .

Next, notice there are three distinct possibilities with respect to subgroups  $N$  and  $M$

- (A)  $N$  and  $M$  are not comparable;
- (B)  $M < N$ ;
- (C)  $N < M$  (for example, the above mentioned example [3]).

## 1. Abelian groups with (A)

In this section we suppose  $M$  and  $N$  are not comparable and  $L(G) = [0, M] \cup [N, G]$ . In this case  $[0, M] \cap [N, G] = \emptyset$  (otherwise  $N \leq M$ ). We list a few straightforward remarks:

- (a)  $M \cap N$  is the largest element in  $[0, N)$  and  $M + N$  is the smallest element in  $(M, G]$ .
- (b)  $L(M + N) = [0, M] \cup [N, N + M]$ , that is,  $N + M$  has property (A).
- (c)  $L(G/(M \cap N)) = [0, M/(M \cap N)] \cup [N/(M \cap N), G/(M \cap N)]$ , that is,  $G/(M \cap N)$  has property (A).
- (d)  $(M + N)/(M \cap N)$  has property (A).

Actually, more can be proved:

**LEMMA 1.1.** *If  $L(G) = [0, M] \cup [N, G]$ , there is a prime number  $p$  such that*

- (a)  $N$  is a (co)cyclic  $p$ -group and  $M \cap N = pN$  is maximal in  $N$ ;
- (b)  $G/M$  and  $G/(M + N)$  are  $p$ -groups.

**PROOF.** (a) We have already noticed that  $N$  has to be cyclic. By the above remark (a),  $N$  is a (co)cyclic  $p$ -group (for a suitable prime number  $p$ ). Moreover, since  $M \cap N$  is its largest (proper) subgroup,  $pN = M \cap N$ .

To prove (b), we observe that  $G/M$  is a cocyclic group since it has a smallest subgroup, namely  $(M+N)/M$ . Moreover, since  $(M+N)/M \cong N/(N \cap M) \cong \mathbb{Z}(p)$ ,  $G/M$  is a cocyclic  $p$ -group, and so  $G/(M+N)$  has the same property.  $\square$

Therefore the subgroup lattice is represented by the diagram shown in Figure 1.




FIGURE 1.

If  $N \simeq \mathbb{Z}(p^k)$  it is readily seen that for  $k = 1$ ,  $N$  is minimal and hence the sum  $N + M$  is direct (otherwise  $N \cap M = N$  and  $N, M$  are comparable). Actually this is the only case  $N \cap M = 0$ .

The following lemma will be used in the proofs of the main results of both this and next sections.

**LEMMA 1.2.** *For a group  $G$  and  $g \in G$ , let  $p$  be a prime such that  $K = G/\langle g \rangle$  is cocyclic  $p$ -group. If  $h_p(g) \neq 0$  and  $G$  is not infinite cyclic, then  $G = H_1 \oplus H_2$  for cocyclic  $p$ -group  $H_1$  and finite cyclic group  $H_2$  of coprime order with  $p$  such that  $H_2 \leq \langle g \rangle$  ( $H_2 = 0$  is not excluded).*

**PROOF.** Since for cocyclic group  $G$  the decomposition is trivial, suppose  $G$  is not cocyclic (and so  $g \neq 0$ ). As  $r(G) \leq r(K) + r(\langle g \rangle) = 2$ , we have  $r(G) = 2$  and by  $r_0(G) = r_0(\langle g \rangle) + r_0(K) \leq 1$ , we obtain  $G = H_1 \oplus H_2$  with  $r(H_1) = r(H_2) = 1$ , that is, each  $H_i$  is cocyclic or infinite cyclic (if  $r_0(G) = 1$ , the torsion subgroup of  $G$  is cocyclic, hence  $G$  splits). If  $g = h_1 + h_2$  with  $h_i \in H_i$ , since  $h_p(g) \geq 1$ , there exist  $x_1 \in H_1$  and  $x_2 \in H_2$  such that  $px_1 = h_1$  and  $px_2 = h_2$ . Moreover,  $L(G/\langle g \rangle)$  is a chain and we can suppose  $x_2 + \langle g \rangle \in (\langle x_1 \rangle + \langle g \rangle)/\langle g \rangle$ .

Thus  $x_2 \in \langle x_1 \rangle + \langle g \rangle$  and  $x_2 = sx_1 + tg$  or  $px_2 = spx_1 + tpg$  for suitable integers  $s, t$ . Hence  $h_2 = sh_1 + tp(h_1 + h_2)$  and, the sum  $H_1 \oplus H_2$  being direct,  $(tp - 1)h_2 = 0$ .

If  $h_2 = 0$  then  $g \in H_1$  and  $K$  is cocyclic if and only if  $\langle g \rangle = H_1$  or  $H_2 = 0$ . In the first case  $h_p(g) = 0$ , hence  $H_2 = 0$  and  $G = H_1$  is a cocyclic  $p$ -group (since, by hypothesis,  $G$  is not infinite cyclic).

If  $h_2 \neq 0$ , the order of  $h_2$  (say  $l$ ) is finite and coprime with  $p$ . Therefore,  $H_2$  is a cocyclic  $q$ -group (if  $l$  is a power of the prime  $q$ ) and this implies  $H_2 \leq \langle g \rangle$  (otherwise  $G/\langle g \rangle$  is not  $p$ -group). Hence there exists a nonzero integer  $k$  such that  $h_2 = kh_1 + kh_2$ , and so  $kh_1 = 0$ . Then  $H_1$  is also cocyclic and necessarily a  $p$ -group.  $\square$

Here is the structure theorem for case (A):

**THEOREM 1.1.** *A group  $G$  satisfies (A) if and only if  $G$  is torsion with a cocyclic primary component and  $r(G) > 1$ .*

**PROOF.** According to Lemma 1.1, let  $p$  be a prime such that  $N = \langle a \rangle$  is cyclic of order  $p^k$ . If  $m \in M \setminus N$  then  $m + a \notin M$  (since  $a \notin M$ ) and  $N \leq \langle m + a \rangle$ . Since  $N \neq 0$  is torsion,  $m + a$  and therefore  $m$  are of finite order. Hence  $M$  and, together with  $G/M$ ,  $G$  are torsion.

Further, we show that  $M_p \subseteq N$ . Indeed, if  $m \in M_p$ , again,  $N \subseteq \langle a + m \rangle$  so that  $a = s(a + m)$  and  $(1 - s)a = sm \in N \cap M = pN$  for a suitable nonzero integer  $s$ . Thus  $s \equiv 1 \pmod{p}$  and let  $t$  be an inverse of  $s$  modulo the order of  $m \in M_p$ . Thus  $m = tsm = t(1 - s)a \in N$ .

Now,  $N$  and  $M$  being not comparable,  $M_p \subset N$  and hence

$$pN = M \cap N = M_p \cap N = M_p.$$

Since  $M_p = pN \leq G_p$ , Lemma 1.2 shows that  $G_p$  is a cocyclic group.

Conversely, suppose  $G = G_p \oplus K$  with  $G_p \simeq \mathbb{Z}(p^l)$ ,  $K \neq 0$ ,  $K_p = 0$  and take  $N = G_p[p] = \langle a \rangle$  and  $M = K$ . If  $H$  is a subgroup of  $G$  such that  $H \not\leq K$  we show  $N \leq H$ .

Indeed, since  $H \not\leq K$ , there is an element  $h \in H \setminus K$ . If this element decomposes as  $h = g_p + k$  ( $g_p \in G_p, k \in K$ ), then  $g_p \neq 0$  and for a suitable multiple  $p^s h = p^s(g_p + k)$  we have  $0 \neq p^s g_p \in N$  respectively  $p^s k \in K$ . Since  $K$  is torsion and  $K_p = 0$ , denoting by  $u$  the order of  $p^s k$ ,  $u$  and  $p$  are coprime and  $up^s g_p \in H$ . Finally,  $p^s g_p \in H$  and thus  $N = \langle p^s g_p \rangle \leq H$ .  $\square$

**REMARKS.** (1) The referee pointed out that a proof in Case (A) can be reduced to the proof of Case (B) using Lemma 1.1. Our proof uses Lemma 1.2 in both cases.

(2) With above notations,  $G/M = \bigoplus_{q \in \mathbb{P}} (G_q/M_q)$  is a  $p$ -group. Hence  $G_q = M_q$  for all primes  $q \neq p$  and  $M = pN \oplus \left( \bigoplus_{q \neq p, q \in \mathbb{P}} G_q \right)$ .




FIGURE 2.

## 2. Abelian groups with (B)

Now we deal with subgroup lattices of the type shown in Figure 2. Here again  $[0, M] \cap [N, G] = \emptyset$ .

Although the following result was already stated in [1], we supply a specific ‘abelian’ proof.

**LEMMA 2.1.**  *$G$  is an abelian BP-group if and only if there is a prime  $p$  and  $k \in \mathbb{N}^* \cup \{\infty\}$ ,  $k \geq 2$  such that  $G \simeq \mathbb{Z}(p^k)$ .*

**PROOF.** If  $L(G) = [0, H] \cup [H, G]$ , then (as noticed in the introduction)  $H$  is a cyclic subgroup. If  $p$  is a prime such that  $pH \neq H$ , then  $H/pH$  is simple, and using again  $L(G) = [0, H] \cup [H, G]$ , it is the smallest nonzero subgroup of  $G/pH$ . Hence  $G/pH$  is cocyclic and, having elements of order  $p$  (in  $H/pH$ ), must be a  $p$ -group. Since an infinite cyclic group is not a BP-group, using Lemma 1.2, we obtain  $G = H_1 \oplus H_2$  with cocyclic  $p$ -group  $H_1$ , cyclic  $q$ -group  $H_2$ ,  $q$  and  $p$  are coprime and  $H_2 \leq pH \leq H$ . Obviously,  $H_1 \not\leq H$  (otherwise  $G = H$ ) so that  $H_2 \leq H \leq H_1$ . This implies  $H_2 = 0$ , and so  $G$  is cocyclic. Since  $\mathbb{Z}(p)$  is not satisfying (B),  $G$  has the requested form.

The converse is immediate (the subgroup lattice of  $\mathbb{Z}(p^n)$  with  $n \in \mathbb{N} \cup \{\infty\}$ ,  $n \geq 2$  is a chain with at least 3 elements). □

Using this we obtain at once

**THEOREM 2.1.** *A group satisfies (B) if and only if  $G \simeq \mathbb{Z}(p^n)$  with  $n \geq 3$ .*

**PROOF.** If  $L(G) = [0, M] \cup [N, G]$  and  $M \leq N$  then  $L(G) = [0, N] \cup [N, G]$




FIGURE 3.

and so  $G$  is a BP-group. Hence  $G$  is cocyclic. Since the conditions  $0 \neq M \neq N \neq G$  require at least 4 elements in  $L(G)$ ,  $G \simeq \mathbb{Z}(p^n)$  with  $n \geq 3$ .

The converse is obvious. □

### 3. Abelian groups with (C)

In this section we consider two proper subgroups  $N < M$  such that  $L(G) = [0, M] \cup [N, G]$ . Thus the subgroup lattice looks as shown in Figure 3.

Now  $L(G) = [0, M] \cup [N, G]$  and  $[0, M] \cap [N, G] = [M, N]$ . Moreover,  $[0, N] \subseteq [0, M]$  and  $[M, G] \subseteq [N, G]$ .

**THEOREM 3.1.** *If a group  $G$  satisfies (C) then  $G$  is a torsion group and there exists a prime  $p$  such that  $G_p$  is a BP-group or satisfies (C). Conversely, if  $G$  is a torsion group,  $G_p \neq G$  for a prime  $p$  and  $G_p$  is a BP-group or satisfies (C), then  $G$  satisfies (C).*

**PROOF.** Let  $0 < N < M < G$  be such that  $L(G) = [0, M] \cup [N, G]$ .

If  $G$  is not a torsion group, there exists an infinite order element  $x \in G$  such that  $x \notin M$  (otherwise, since the infinite order elements generate any group,  $M = G$ ). Then  $0 < N \leq M \cap \langle x \rangle < \langle x \rangle$ . If  $L \leq \langle x \rangle$  then  $L \leq M$  or  $N \leq L$ , hence  $L \leq M \cap \langle x \rangle$  or  $N \leq L$ . Therefore  $\langle x \rangle$  is a BP-group or satisfies (C), but it is easy to see that no infinite cyclic group satisfies these properties (as for (C), if  $0 < n\mathbb{Z} < m\mathbb{Z} < \mathbb{Z}$  and  $p$  is a prime not dividing  $n$ , then  $p\mathbb{Z} \notin [0, m\mathbb{Z}] \cup [n\mathbb{Z}, \mathbb{Z}]$ ). This contradiction shows that  $G$  is a torsion group.

Suppose no component  $G_p$  is a BP-group or satisfies (C). Since  $M \neq G$ , there exists a prime  $p$  such that  $M_p \neq G_p$ . If  $N_p = 0$ , then  $G_p \subseteq M$  ( $N \subseteq G_p$  is not possible,  $N$  being a proper subgroup), hence  $M_p = G_p$ . Therefore  $0 < N_p \leq M_p < G_p$  and  $L(G_p) \neq [0, M_p] \cup [N_p, G_p]$ . Then we can find  $H_p \leq G_p$  such that  $H_p \setminus M_p \neq \emptyset$  and  $N_p \setminus H_p \neq \emptyset$ . It follows  $H_p \setminus M \neq \emptyset$  and  $N \setminus H_p \neq \emptyset$ , a contradiction.

Conversely, suppose  $G$  is torsion and  $G_p$  is a BP-group or satisfies (C). Then we can find subgroups  $0 < N_p \leq M_p < G_p$  such that  $L(G_p) = [0, M_p] \cup [N_p, G_p]$ . Set  $M = M_p \oplus (\bigoplus_{q \neq p} G_q)$  and  $N = N_p$ . Thus  $0 < N < M < G$ . If  $H \leq G$ , then  $H = H_p \oplus (\bigoplus_{q \neq p} H_q)$  with  $H_p \leq G_p$  and  $\bigoplus_{q \neq p} H_q \leq \bigoplus_{q \neq p} G_q$ . If  $N_p \leq H_p$ , then  $H \in [N, G]$  and if  $H_p \leq M_p$ , then  $H \leq M_p \oplus (\bigoplus_{q \neq p} H_q) \leq M_p \oplus (\bigoplus_{q \neq p} G_q) = M$ .

Actually,  $G_p \neq G$  is needed only for a BP-group  $G_p$  not satisfying (C).  $\square$

**THEOREM 3.2.** *A  $p$ -group  $G$  satisfies (C) if and only if  $G \cong \mathbb{Z}(p^n) \oplus B$  such that (i)  $B \neq 0$ ,  $n \in \mathbb{N}^* \cup \{\infty\}$  and  $p^l B = 0$  holds for a positive integer  $l < n$ , or (ii)  $B = 0$  and  $n > 2$ .*

**PROOF.** If  $G$  satisfies (C), we can suppose  $N = \langle a \rangle \cong \mathbb{Z}(p)$ . Let  $l > 0$  be the smallest positive integer such that there exists  $x \in G \setminus M$  with  $p^l x = a$ . Let  $b \in G[p] \setminus \langle a \rangle$  and suppose  $h_p(b) \geq l$ . Then  $b = p^l y$  for some  $y \in M$  (if  $y \notin M$  we have  $a \in \langle y \rangle$ , hence the rank of  $\langle y \rangle[p]$  is at least 2, a contradiction). Thus  $x + y \notin M$ , and there exists a positive integer  $k$  such that  $kx + ky = a$ . If  $k = p^r m$  with  $\gcd(m; p) = 1$  then  $p^r(mx + my) = a$ , hence  $l \leq r$ . Moreover,  $l \leq r$  implies  $ky \in \langle a \rangle$  and  $a \in \langle y \rangle$  follows, a contradiction. Then  $h_p(b) < l$  for all  $b \in G[p] \setminus \langle a \rangle$  and so  $p^l G[p] = \langle a \rangle$ . Hence  $p^l G$  is a cocyclic group.

If  $p^l G$  is a cyclic group then  $G$  is bounded and (using [2, 27.2])  $G = H \oplus B$  where  $H \cong \mathbb{Z}(p^n)$  with  $n \geq l + 1$ ,  $a \in H$  and  $p^l B = 0$  (otherwise there is  $b \in B[p]$  with  $h_p(b) \geq l$ ). If  $p^l G$  is a quasicyclic group, then  $G = p^l G \oplus B$  and  $p^l B = 0$ .

Moreover, if  $B = 0$  then  $G \cong \mathbb{Z}(p^n)$  and condition  $M \neq N$  implies  $n > 2$ .

Conversely, if  $B = 0$  then  $G$  satisfies condition (C) for  $N = p^{n-1}G$  and  $M = pG$ . If  $B \neq 0$  we choose  $G = H \oplus B$  with  $H \simeq \mathbb{Z}(p^n)$ ,  $0 < l < n$  such that  $p^l B = 0$ ,  $N = H[p] = \langle a \rangle \cong \mathbb{Z}(p)$  and  $M = A + B$  where  $A$  is the subgroup of  $H$  of order  $p^l$  (obviously containing  $N$  - the subgroup lattice of  $H$  being a chain with a smallest element). If  $X$  is a subgroup of  $G$  such that  $X \not\subseteq [0, M]$ , then there exists  $x = h + b \in X \setminus M$  with  $h \in H$  and  $b \in B$  such that  $p^r = \text{ord}(h) > p^l$  (otherwise  $h \in A$  and  $x \in M$ ). By  $p^l B = 0$  hypothesis,  $0 \neq p^{r-1}x = p^{r-1}h \in H[p] = N$ , hence  $\langle p^{r-1}h \rangle = N$  is included in  $X$ .  $\square$

The only BP-groups which do not satisfy (B), nor (C) are  $\mathbb{Z}(p^2)$  for any prime number  $p$ . Hence




FIGURE 4.

**COROLLARY 3.1.** *A group  $G$  satisfies (C) if and only if it is a torsion group with a primary component  $\mathbb{Z}(p^n)$  for  $n \geq 3$ , or  $G_p \cong \mathbb{Z}(p^n) \oplus B$  with  $n > 1$  or  $\infty$  and  $p^l B = 0$  holds for a nonnegative integer  $l < n$ .*

### 4. Comments

1. There are groups satisfying both conditions (A) and (C). As an example take

$$G = \mathbb{Z}(12) = \langle a, b \mid 3a = 4b = 0 \rangle.$$

Denoting by  $N = \langle a \rangle$  and  $M = \langle b \rangle$  the subgroup lattice looks as shown in Figure 4 (a). Thus  $L(G) = [0, M] \cup [N, G]$  for (A), and  $L(G) = [0, N + 2M] \cup [2M, G]$  for (C).

2. If a group  $G$  satisfies, say, the condition (C) the pair  $M, N$  of subgroups is not necessarily unique. As an example, take the group

$$G = \mathbb{Z}(2) \oplus \mathbb{Z}(8) = \langle a, b \mid 2a = 8b = 0 \rangle.$$

If we denote by  $N = \langle a \rangle$ ,  $M = \langle b \rangle$ ,  $S = \langle a + 2b \rangle$ ,  $T = \langle a + b \rangle$ ,  $U = \langle a + 4b \rangle$ , the subgroup lattice is now as shown in Figure 4 (b) and

$$L(G) = [0, N + 2M] \cup [2M, G] = [0, N + 2M] \cup [4M, G].$$

3. Our results generalize to lattices with 0 and 1, more or less arbitrary. In what follows we state some of these lattice versions.

- If a lattice  $L$  satisfies condition (A), that is,  $L = [0, m] \cup [n, 1]$  with incomparable elements  $m, n$  then






FIGURE 5.

- (a)  $[0, m \vee n] = [0, m] \cup [n, m \vee n]$  that is,  $[0, m \vee n]$  satisfies condition (A);
- (b)  $[m \wedge n, 1] = [m \wedge n, m] \cup [n, 1]$  that is,  $[m \wedge n, 1]$  satisfies condition (A);
- (c)  $[m \wedge n, m \vee n]$  satisfies condition (A).

- Every direct product of two lattices, the first being a finite chain and the second having 0 and 1, satisfies condition (A).

**PROOF.** One uses the diagram shown in Figure 5 (for the sake of simplicity we have considered a chain with only two elements).

Denoting the chain by  $\{a, b\}$  and using elements in the Cartesian product  $\{a, b\} \times L$ , decomposition in the required intervals is  $[(a, 0), (a, 1)] \cup [(b, 0), (b, 1)]$ . □

A family of torsion groups is said to be *coprime* if the orders of elements in any two members are coprime. Using an early Theorem of Suzuki (see [3]): *the groups with decomposable subgroup lattices are exactly the direct sums of coprime groups*, we have an alternative proof for sufficiency of Theorem 1.1 in the special case  $k = 1$ .

**PROOF FOR SUFFICIENCY OF THEOREM 1.1 ( $k = 1$ ).** Let  $G$  be a torsion group of rank  $r(G) > 1$  with a simple  $p$ -component, that is  $G = N \oplus M$  with  $|N| = p$  and  $M_p = 0$ . Thus  $N$  and  $M$  are coprime,  $L(G) \simeq L(N) \times L(M)$  and  $L(N)$  is a chain with two elements. Applying the previous result,  $L(G)$  satisfies condition (A). □

- Complemented lattices are not satisfying condition (C).
- Let  $\{L_i, i \in I\}$  be an arbitrary set of bounded (that is, with  $0_i$  and  $1_i$ ) lattices, at least one of these satisfying condition (C). Then the direct product  $L = \prod_{i \in I} L_i$  satisfies condition (C). Conversely, if  $L$  satisfies condition (C), that is,  $L = [0, \alpha] \cup [\beta, 1]$  and for an index  $j \in I, 0_j < \beta_j < \alpha_j < 1_j$ , then  $L_j$  satisfies condition (C).

- If a lattice satisfies condition (C), that is,  $L = [0, m] \cup [n, 1]$ , then  $m$  is essential and  $n$  is superfluous in  $L$ . Moreover, every element disjoint with  $n$  belongs to  $[0, m]$ .

Finally we mention the lattice version of our initial proof of case (A):

- Let  $L$  be a modular lattice,  $n$  an atom and  $m$  a dual atom in  $L$  such that  $1 = n \vee m$  and  $n \wedge m = 0$ . Then  $L = [0, m] \cup [n, 1]$  if and only if for every element  $v$  in  $[0, m]$ ,  $n$  has a unique (relative) complement (namely  $v$ ) in the sublattice  $[0, n \vee v]$ .

Using this, one can show that, excepting the case  $1 = n \vee m$  and  $n \wedge m = 0$ , (C) follows from (A).

### Acknowledgement

Thanks are due to the referee for his (her) valuable suggestions and improvements.

### References

- [1] G. Călugăreanu and M. Deaconescu, 'Breaking points in subgroup lattices', in: *Proceedings of Groups St Andrews 2001 in Oxford* (Cambridge University Press, Cambridge, 2003) pp. 59–62.
- [2] L. Fuchs, *Infinite abelian groups*, Vol. 1, 2 (Academic Press, New York, 1970, 1973).
- [3] R. Schmidt, *Subgroup lattices of groups*, Expositions in Mathematics 14 (Walter de Gruyter, 1994).
- [4] J. Tuma, 'Intervals in subgroup lattices of infinite groups', *J. Algebra* (2) **125** (1989), 367–399.
- [5] P. M. Whitman, 'Lattices, equivalence relations, and subgroups', *Bull. Amer. Math. Soc.* **52** (1946), 507–522.

Faculty of Mathematics  
and Informatics  
Babes-Bolyai University  
Cluj-Napoca  
Romania  
e-mail: [bodo@math.ubbcluj.ro](mailto:bodo@math.ubbcluj.ro)

Dept. Mathematics and Computer Sci  
Faculty of Science  
Kuwait University  
State of Kuwait  
e-mail: [calu@mcs.sci.kuniv.edu.kw](mailto:calu@mcs.sci.kuniv.edu.kw)